

Sugar Mover



SPECIMEN

3-0-0 GUARANTEED ANALYSIS

Total Nitrogen (N).....	3.0%
3% Water Soluble Nitrogen	
Boron (B).....	8.0%
Molybdenum (Mo).....	0.004%

Derived from boron ethanolamine and sodium molybdate.

F2399

Z-SUGARM
Rev: 21C10

Density: 10.6 lb/gal (1.27 kg/L) @ 68°F (20°C)

NET CONTENT	NET WEIGHT
<input type="checkbox"/> 2.50 gal (9.5 L)	26.5 lb (12.0 kg)
<input type="checkbox"/> 2.64 gal (10.0 L)	28.0 lb (12.7 kg)
<input type="checkbox"/> 5.00 gal (18.9 L)	53.0 lb (24.1 kg)
<input type="checkbox"/> 5.28 gal (20.0 L)	56.0 lb (25.4 kg)
<input type="checkbox"/> 55.00 gal (208.2 L)	583.0 lb (264.8 kg)
<input type="checkbox"/> 275.00 gal (1041.0 L)	2915.0 lb (1324.1 kg)
<input type="checkbox"/> BULK	



WARNING

KEEP OUT OF REACH OF CHILDREN; MAY BE HARMFUL IF SWALLOWED, INHALED OR IN CONTACT WITH SKIN; MAY CAUSE EYE IRRITATION. MAY CAUSE DAMAGE THROUGH PROLONGED EXPOSURE.

PRECAUTIONS:

Do not breathe dust/fume/gas/mist/vapors/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in well-ventilated area. Avoid release to environment. Wear protective gloves/protective clothing/eye protection/face protection. Keep container tightly closed. Keep cool and protected from sunlight.

If medical advice is needed, have product container or label at hand. For more details regarding the safe use/handling, storage, and disposal of this product as well as additional hazards and first aid measures, please obtain the current Safety Data Sheet at www.stollerusa.com.

STORAGE AND DISPOSAL

We recommend storing pails a maximum of three pails high. Do not stack pallets. Store container tightly closed in a cool/well-ventilated dry place protected from sunlight above 40°F. Do not allow container to freeze. Store locked up. DISPOSE of contents/container at a permitted facility or as advised by your local regulatory authority. Do not contaminate water, food or feed by storage or disposal. Do not apply directly to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water. Do not reuse empty container.

WARNING: This product contains BORON at a level that may be detrimental to growing plants. Use of boron on crops other than those recommended may result in serious injury to those crops.

WARNING: The application of fertilizing materials containing MOLYBDENUM may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

Always read and follow label instructions.

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | www.stollerusa.com | 800-539-5283 or 713-461-1493 | ©2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.

GENERAL INFORMATION

SUGAR MOVER is a soil or foliar applied source of soluble boron with the added benefit of molybdenum and is non-phytotoxic when used as directed.

DIRECTIONS FOR USE

It may be a violation of federal law to use this product inconsistent with its labeling.

SUGAR MOVER can be applied to the plant foliage by spraying, or to the root zone by flood, drip, overhead, pivot and micro irrigation systems. Apply the recommended amount in a volume sufficient to provide uniform coverage.

RECOMMENDED USES

FOR FERTIGATION: Apply **SUGAR MOVER** through irrigation system to the root zone using 1 to 4 pints per planted acre. Repeat applications can be made every 7-21 days as needed. **SUGAR MOVER** can be incorporated into liquid fertilizer or applied separately; DO NOT blend into fertilizers intended for in furrow or application to seed.

FOR NITROGEN SIDE DRESS APPLICATION: Apply **SUGAR MOVER** at 1 quart per 100 units of actual nitrogen. Can be blended into liquid 28% or 32% UAN or impregnated onto dry urea or AMS.

FOR FOLIAR APPLICATION: Apply **SUGAR MOVER** to the plant foliage by spraying 1 to 4 pints per planted acre. Repeat applications can be made every 7-21 days as needed. Do not exceed 4 pints per acre per foliar application.

Use **SUGAR MOVER** at the recommended rate on the following crops:

FIELD CROPS: Alfalfa, bean crops, canola, clover, corn, cotton, flax, forage grasses, peanut, potato, rape (canola), rice, sorghum, soybean, sugar beet, sugarcane, sunflower, tobacco, and wheat, barley, oat, or other cereal grain.

VEGETABLE CROPS: Artichoke, asparagus, broccoli, cauliflower, cabbage, kale, and other cruciferous vegetables; leaf lettuce, head lettuce, dandelion, radicchio, and other leafy vegetables; cucumber, melon, pumpkin, squash, and other cucurbits, onion, garlic, and other bulb vegetables; tomato, pepper, eggplant and other fruiting vegetables; strawberry; carrot, sweet potato, radish, beet and other root and tuber vegetables.

FRUIT AND NUT CROPS: Apple, avocado, blueberry and other bush berries, blackberry, raspberry and other cane berries, sweet orange, lemon, mandarin and other citrus, date palm, fig, grapes, hops, oil palm, olive, persimmon, cherry, peach, plums, prune, nectarine, apricot and other stone fruit, almond, pecan, pistachio, walnut and other tree nuts, pear, pomegranate, and subtropical fruit.

HARVEST PREPARATION APPLICATION: Use as directed on the following: **Alfalfa:** Apply 1 quart 8 days before cutting. **Asparagus:** 1 to 2 quarts per acre (foliar) 3 weeks before cutting fern. **Canning Tomato:** Apply 1 quart per acre 14 days before harvest. **Canola:** Apply 1 quart per acre at flowering before pods appear. **Corn:** Apply 1 to 2 quarts per acre 15 days before tasseling. **Cotton:** Apply 1 to 2 quarts per acre 2 ½ to 3 weeks before defoliation. **Onion/Garlic/Carrot:** Apply 1 to 2 quarts per acre 2 ½ to 3 weeks before harvest. **Potato:** 2 quarts/acre (foliar) 2 ½ to 3 weeks before killing; if vines are not killed, apply 3 weeks before harvest. **Soybean:** Apply 1 quart per acre at bloom before pod emergence. **Sugar Beet:** Apply 2 quarts per acre when plants should normally be senescing. **Wheat:** Apply 1 quart per acre 15 days before the seed heads appear. **Fruit and Nut Crops: Apple, Cherry and Peach:** Apply 1 pint per acre 14 days, 7 days and 4 days before harvest. **Strawberry Nursery Daughter Plant Production:** Apply 1 to 2 quarts per acre, two times, 10 to 14 days apart, beginning 4 to 5 weeks before harvest.

POSTHARVEST/HIBERNATION APPLICATION ON

PERENNIALS: Apply single or multiple treatments of 1 to 4 pints of **SUGAR MOVER** per acre after harvest and before leaf fall. A post-harvest application can prepare trees (i.e., almonds and other tree fruit), vines and bushes (i.e., perennial berry crops) and range crops (alfalfa, grasses and other perennial forages) for dormancy, bud-break and flowering the following season.

MIXING INSTRUCTIONS: Follow this mixing order (1) Water (2) **SUGAR MOVER** (3) Pesticide. **SUGAR MOVER** will disperse in water with little agitation. **SUGAR MOVER** is compatible with most fertilizers and pesticides. Always conduct a jar test when using new or untried combinations.

Notice: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller makes any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Always read and follow label instructions.

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | www.stollerusa.com | 800-539-5283 or 713-461-1493 | ©2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.